

A Crash Course on Discrete Probability

Events and Probability

Consider a random process (e.g., throw a die, pick a card from a deck)

- Each possible outcome is a **simple event** (or sample point).
- The **sample space** Ω is the set of all possible simple events.
- An **event** is a set of simple events (a subset of the sample space).
- With each simple event E we associate a real number $0 \leq \Pr(E) \leq 1$ which is the **probability** of E .

Probability Space

Definition

A **probability space** has three components:

- 1 A **sample space** Ω , which is the set of all possible outcomes of the random process modeled by the probability space;
- 2 A family of sets \mathcal{F} representing the allowable **events**, where each set in \mathcal{F} is a subset of the sample space Ω ;
- 3 A **probability function** $\Pr : \mathcal{F} \rightarrow \mathbf{R}$, satisfying the definition below.

In a **discrete** probability space we use $\mathcal{F} =$ “all the subsets of Ω ”

Probability Function

Definition

A **probability function** is any function $\Pr : \mathcal{F} \rightarrow \mathbf{R}$ that satisfies the following conditions:

- 1 For any event E , $0 \leq \Pr(E) \leq 1$;
- 2 $\Pr(\Omega) = 1$;
- 3 For any finite or countably infinite sequence of **pairwise mutually disjoint** events E_1, E_2, E_3, \dots

$$\Pr \left(\bigcup_{i \geq 1} E_i \right) = \sum_{i \geq 1} \Pr(E_i).$$

The probability of an event is the sum of the probabilities of its simple events.

Examples:

Consider the random process defined by the outcome of rolling a die.

$$\mathcal{S} = \{1, 2, 3, 4, 5, 6\}$$

We assume that all “facets” have equal probability, thus

$$\Pr(1) = \Pr(2) = \dots \Pr(6) = 1/6.$$

The probability of the event “odd outcome”

$$= \Pr(\{1, 3, 5\}) = 1/2$$

Assume that we roll two dice:

$\mathcal{S} =$ all ordered pairs $\{(i, j), 1 \leq i, j \leq 6\}$.

We assume that each (ordered) combination has probability $1/36$.

Probability of the event “sum = 2”

$$\Pr(\{(1, 1)\}) = 1/36.$$

Probability of the event “sum = 3”

$$\Pr(\{(1, 2), (2, 1)\}) = 2/36.$$

Let $E_1 =$ “sum bounded by 6”,

$$E_1 = \{(1, 1), (1, 2), (1, 3), (1, 4), (1, 5), (2, 1), (2, 2), \\ (2, 3), (2, 4), (3, 1), (3, 2), (3, 3), (4, 1), (4, 2), (5, 1)\}$$

$$\Pr(E_1) = 15/36$$

Let $E_2 =$ “both dice have odd numbers”, $\Pr(E_2) = 1/4$.

$$\Pr(E_1 \cap E_2) =$$

$$\Pr(\{(1, 1), (1, 3), (1, 5), (3, 1), (3, 3), (5, 1)\}) =$$

$$6/36 = 1/6.$$

Conditional Probability

What is the probability that a random student at University of Ioannina was born in Ioannina.

E_1 = the event “born in Ioannina.”

E_2 = the event “a student in Uoi.”

The conditional probability that a student at Uoi was born in Ioannina is written:

$$\Pr(E_1 | E_2).$$

Computing Conditional Probabilities

Definition

The **conditional probability** that event E occurs given that event F occurs is

$$\Pr(E | F) = \frac{\Pr(E \cap F)}{\Pr(F)}.$$

The conditional probability is only well-defined if $\Pr(F) > 0$.

By conditioning on F we restrict the sample space to the set F . Thus we are interested in $\Pr(E \cap F)$ “normalized” by $\Pr(F)$.

Example

What is the probability that in rolling two dice the sum is 8 given that the sum was even?

Example

What is the probability that in rolling two dice the sum is 8 given that the sum was even?

$E_1 =$ “sum is 8” ,

$E_2 =$ “sum even” ,

Example

What is the probability that in rolling two dice the sum is 8 given that the sum was even?

$E_1 =$ “sum is 8” ,

$E_2 =$ “sum even” ,

$$\Pr(E_1) = \Pr(\{(2, 6), (3, 5), (4, 4), (5, 3), (6, 2)\}) = 5/36$$

Example

What is the probability that in rolling two dice the sum is 8 given that the sum was even?

$$E_1 = \text{"sum is 8"},$$

$$E_2 = \text{"sum even"},$$

$$\Pr(E_1) = \Pr(\{(2, 6), (3, 5), (4, 4), (5, 3), (6, 2)\}) = 5/36$$

$$\Pr(E_2) = 1/2 = 18/36.$$

Example

What is the probability that in rolling two dice the sum is 8 given that the sum was even?

$E_1 =$ “sum is 8”,

$E_2 =$ “sum even”,

$$\Pr(E_1) = \Pr(\{(2, 6), (3, 5), (4, 4), (5, 3), (6, 2)\}) = 5/36$$

$$\Pr(E_2) = 1/2 = 18/36.$$

$$\Pr(E_1 | E_2) = \frac{\Pr(E_1 \cap E_2)}{\Pr(E_2)} = \frac{5/36}{1/2} = 5/18.$$

A Useful Identity

Assume two events A and B .

$$\begin{aligned}\Pr(A) &= \Pr(A \cap B) + \Pr(A \cap B^c) \\ &= \Pr(A \mid B) \cdot \Pr(B) + \Pr(A \mid B^c) \cdot \Pr(B^c)\end{aligned}$$

A Useful Identity

Assume two events A and B .

$$\begin{aligned}\Pr(A) &= \Pr(A \cap B) + \Pr(A \cap B^c) \\ &= \Pr(A \mid B) \cdot \Pr(B) + \Pr(A \mid B^c) \cdot \Pr(B^c)\end{aligned}$$

Example:

What is the probability that a random person has height > 1.75 ?

We choose a random person and let A the event that “the person has height > 1.75 .”

We want $\Pr(A)$.

A Useful Identity

Assume two events A and B .

$$\begin{aligned}\Pr(A) &= \Pr(A \cap B) + \Pr(A \cap B^c) \\ &= \Pr(A \mid B) \cdot \Pr(B) + \Pr(A \mid B^c) \cdot \Pr(B^c)\end{aligned}$$

Example:

What is the probability that a random person has height > 1.75 ?

We choose a random person and let A the event that “the person has height > 1.75 .”

We want $\Pr(A)$.

Assume we know that the probability that a man has height > 1.75 is 54% and that a woman has height > 1.75 is 4%.

A Useful Identity

Assume two events A and B .

$$\begin{aligned}\Pr(A) &= \Pr(A \cap B) + \Pr(A \cap B^c) \\ &= \Pr(A \mid B) \cdot \Pr(B) + \Pr(A \mid B^c) \cdot \Pr(B^c)\end{aligned}$$

Example:

What is the probability that a random person has height > 1.75 ?

We choose a random person and let A the event that “the person has height > 1.75 .”

We want $\Pr(A)$.

Assume we know that the probability that a man has height > 1.75 is 54% and that a woman has height > 1.75 is 4%.

Define the event B that “the random person is a man.”

Example - a posteriori probability

We are given 2 coins:

- one is a fair coin A
- the other coin, B , has head on both sides

We choose a coin at random, i.e. each coin is chosen with probability $1/2$. We then flip the coin.

Given that we got head, what is the probability that we chose the fair coin A ???

Define a sample space of ordered pairs (*coin*, *outcome*).
The sample space has three points

$$\{(A, h), (A, t), (B, h)\}$$

$$\Pr((A, h)) = \Pr((A, t)) = 1/4$$

$$\Pr((B, h)) = 1/2$$

Define two events:

E_1 = "Chose coin *A*".

E_2 = "Outcome is head".

$$\Pr(E_1 | E_2) = \frac{\Pr(E_1 \cap E_2)}{\Pr(E_2)} = \frac{1/4}{1/4 + 1/2} = 1/3.$$

Bayes Rule

Another way to compute the same thing:

$$\begin{aligned}\Pr(E_1 | E_2) &= \frac{\Pr(E_1 \cap E_2)}{\Pr(E_2)} \\ &= \frac{\Pr(E_2 | E_1) \cdot \Pr(E_1)}{\Pr(E_2)} \\ &= \frac{\Pr(E_2 | E_1) \cdot \Pr(E_1)}{\Pr(E_2 | E_1)\Pr(E_1) + \Pr(E_2 | \overline{E_1})\Pr(\overline{E_1})} \\ &= \frac{1/2 \cdot 1/2}{1/2 \cdot 1/2 + 1 \cdot 1/2} = 1/3.\end{aligned}$$

Independent Events

Definition

Two events E and F are **independent** if and only if

$$\Pr(E \cap F) = \Pr(E) \cdot \Pr(F).$$

Equivalently we can write:

$$\Pr(A | B) = \frac{\Pr(A \cap B)}{\Pr(B)} = \Pr(A).$$

Independent Events, examples

Example: You pick a card from a deck.

- $E =$ "Pick an ace"
- $F =$ "Pick a heart"

Example: You roll a die

- $E =$ "number is even"
- $F =$ "number is ≤ 4 "

Basically, two events are independent if when one happens it doesn't tell you anything about if the other happened.

Complement

If event E has probability $\Pr(E)$, then the complement of the event \bar{E} has probability $1 - \Pr(E)$.

Sometimes it is easier to compute this probability. For example:

$E =$ "In 3 rolls of the dice I get at least one 6"

Computing all combinations of events where this is true is complex. What is the complement of E ?

Complement

If event E has probability $\Pr(E)$, then the complement of the event \bar{E} has probability $1 - \Pr(E)$.

Sometimes it is easier to compute this probability. For example:

$E =$ "In 3 rolls of the dice I get at least one 6"

Computing all combinations of events where this is true is complex. What is the complement of E ?

$\bar{E} =$ "In 3 rolls of the dice I **never** get a 6"

Complement

If event E has probability $\Pr(E)$, then the complement of the event \bar{E} has probability $1 - \Pr(E)$.

Sometimes it is easier to compute this probability. For example:

E = "In 3 rolls of the dice I get at least one 6"

Computing all combinations of events where this is true is complex. What is the complement of E ?

\bar{E} = "In 3 rolls of the dice I **never** get a 6"

$$\Pr(\bar{E}) = (1 - 1/6) \cdot (1 - 1/6) \cdot (1 - 1/6)$$

Complement

If event E has probability $\Pr(E)$, then the complement of the event \bar{E} has probability $1 - \Pr(E)$.

Sometimes it is easier to compute this probability. For example:

E = "In 3 rolls of the dice I get at least one 6"

Computing all combinations of events where this is true is complex. What is the complement of E ?

\bar{E} = "In 3 rolls of the dice I **never** get a 6"

$$\Pr(\bar{E}) = (1 - 1/6) \cdot (1 - 1/6) \cdot (1 - 1/6)$$

$$\Pr(E) = 1 - \Pr(\bar{E}) = 1 - (5/6)^3$$

Random Variable

Definition

A **random variable** X on a sample space Ω is a function on Ω ; that is, $X : \Omega \rightarrow \mathcal{R}$.

A **discrete random variable** is a random variable that takes on only a finite or countably infinite number of values.

Examples:

In practice, a random variable is some random quantity that we are interested in:

- 1 I roll a die, $X = \text{"result"}$

Examples:

In practice, a random variable is some random quantity that we are interested in:

- ① I roll a die, $X = \text{"result"}$
- ② I roll 2 dice, $X = \text{"sum of the two values"}$

Examples:

In practice, a random variable is some random quantity that we are interested in:

- ① I roll a die, $X = \text{"result"}$
- ② I roll 2 dice, $X = \text{"sum of the two values"}$
- ③ Consider a gambling game in which a player flips two coins, if he gets heads in both coins he wins \$3, else he loses \$1. The payoff of the game is a random variable.

Examples:

In practice, a random variable is some random quantity that we are interested in:

- ① I roll a die, $X = \text{"result"}$
- ② I roll 2 dice, $X = \text{"sum of the two values"}$
- ③ Consider a gambling game in which a player flips two coins, if he gets heads in both coins he wins \$3, else he loses \$1. The payoff of the game is a random variable.
- ④ I pick a card, $X = \begin{cases} 1, & \text{if card is an Ace} \\ 0, & \text{otherwise} \end{cases}$

Examples:

In practice, a random variable is some random quantity that we are interested in:

- ① I roll a die, $X = \text{"result"}$
- ② I roll 2 dice, $X = \text{"sum of the two values"}$
- ③ Consider a gambling game in which a player flips two coins, if he gets heads in both coins he wins \$3, else he loses \$1. The payoff of the game is a random variable.
- ④ I pick a card, $X = \begin{cases} 1, & \text{if card is an Ace} \\ 0, & \text{otherwise} \end{cases}$
- ⑤ I pick a random Greek citizen, $X = \text{"weight"}$

Examples:

In practice, a random variable is some random quantity that we are interested in:

- 1 I roll a die, $X = \text{"result"}$
- 2 I roll 2 dice, $X = \text{"sum of the two values"}$
- 3 Consider a gambling game in which a player flips two coins, if he gets heads in both coins he wins \$3, else he loses \$1. The payoff of the game is a random variable.
- 4 I pick a card, $X = \begin{cases} 1, & \text{if card is an Ace} \\ 0, & \text{otherwise} \end{cases}$
- 5 I pick a random Greek citizen, $X = \text{"weight"}$
- 6 I pick 10 random students, $X = \text{"average weight"}$

Examples:

In practice, a random variable is some random quantity that we are interested in:

- 1 I roll a die, $X = \text{"result"}$
- 2 I roll 2 dice, $X = \text{"sum of the two values"}$
- 3 Consider a gambling game in which a player flips two coins, if he gets heads in both coins he wins \$3, else he loses \$1. The payoff of the game is a random variable.
- 4 I pick a card, $X = \begin{cases} 1, & \text{if card is an Ace} \\ 0, & \text{otherwise} \end{cases}$
- 5 I pick a random Greek citizen, $X = \text{"weight"}$
- 6 I pick 10 random students, $X = \text{"average weight"}$
- 7 $X = \text{"Running time of quicksort"}$

Independent random variables

Definition

Two random variables X and Y are **independent** if and only if

$$\Pr((X = x) \cap (Y = y)) = \Pr(X = x) \cdot \Pr(Y = y)$$

for all values x and y .

Independent random variables

- A player rolls 5 dice. The sum in the first 3 dice and the sum in the last 2 dice are independent.

Independent random variables

- A player rolls 5 dice. The sum in the first 3 dice and the sum in the last 2 dice are independent.
- I pick a random card from a deck. The **value** that I got and the **suit** that I got are independent.

Independent random variables

- A player rolls 5 dice. The sum in the first 3 dice and the sum in the last 2 dice are independent.
- I pick a random card from a deck. The **value** that I got and the **suit** that I got are independent.
- I pick a random person in Ioannina. The **age** and the **weight** are **not** independent.

Expectation

Definition

The **expectation** of a discrete random variable X , denoted by $\mathbf{E}[X]$, is given by

$$\mathbf{E}[X] = \sum_i i \Pr(X = i),$$

where the summation is over all values in the range of X .

Think of the expectation as the mean value you would get if you took many, many values of the random variable.

Examples:

- The expected value of one die roll is:

$$E[X] = \sum_{i=1}^6 i \Pr(X = i) = \sum_{i=1}^6 i \frac{1}{6} = 3\frac{1}{2}.$$

- The expectation of the random variable X representing the sum of two dice is

$$E[X] = \frac{1}{36} \cdot 2 + \frac{2}{36} \cdot 3 + \frac{3}{36} \cdot 4 + \dots + \frac{1}{36} \cdot 12 = 7.$$

- Let X take on the value 2^i with probability $1/2^i$ for $i = 1, 2, \dots$

$$E[X] = \sum_{i=1}^{\infty} \frac{1}{2^i} 2^i = \sum_{i=1}^{\infty} 1 = \infty.$$

Linearity of Expectation

Theorem

For any two random variables X and Y

$$E[X + Y] = E[X] + E[Y].$$

Theorem

For any constant c and discrete random variable X ,

$$E[cX] = cE[X].$$

Note: X and Y do not have to be independent.

Examples:

- The expectation of the sum of n dice is . . .

Examples:

- The expectation of the sum of n dice is . . .
- The expectation of the outcome of one die plus twice the outcome of a second die is . . .

Bernoulli Random Variable

A **Bernoulli** or an **indicator** random variable:

$$Y = \begin{cases} 1 & \text{if the experiment succeeds,} \\ 0 & \text{otherwise.} \end{cases}$$

$$\mathbf{E}[Y] = p \cdot 1 + (1 - p) \cdot 0 = p = \mathbf{Pr}(Y = 1).$$

Binomial Random Variable

Assume that we repeat n independent Bernoulli trials that have probability p .

Examples:

- I flip n coins, $X_i = 1$, if the i th flip is “head” ($p = 1/2$)
- I roll n dice, $X_i = 1$, if the i th die roll is a 4 ($p = 1/6$)
- I choose n cards, $X_i = 1$, if the i th card is a J, Q, K ($p = 12/52$.)

Let $X = \sum_{i=1}^n X_i$.

X is a Binomial random variable.

Binomial Random Variable

Definition

A binomial random variable X with parameters n and p , denoted by $B(n, p)$, is defined by the following probability distribution on $j = 0, 1, 2, \dots, n$:

$$\Pr(X = j) = \binom{n}{j} p^j (1 - p)^{n-j}.$$

$\binom{n}{k} = \frac{n!}{k!(n-k)!}$ is the number of ways that we can select k elements out of n .

Expectation of a Binomial Random Variable

$$\begin{aligned}\mathbf{E}[X] &= \sum_{j=0}^n j \Pr(X = j) \\ &= \sum_{j=0}^n j \binom{n}{j} p^j (1-p)^{n-j}\end{aligned}$$

Expectation of a Binomial Random Variable

$$\begin{aligned}\mathbf{E}[X] &= \sum_{j=0}^n j \Pr(X = j) \\ &= \sum_{j=0}^n j \binom{n}{j} p^j (1-p)^{n-j} \\ &= \sum_{j=0}^n j \frac{n!}{j!(n-j)!} p^j (1-p)^{n-j} \\ &= \sum_{j=1}^n \frac{n!}{(j-1)!(n-j)!} p^j (1-p)^{n-j} \\ &= np \sum_{j=1}^n \frac{(n-1)!}{(j-1)!((n-1)-(j-1))!} p^{j-1} (1-p)^{(n-1)-(j-1)} \\ &= np \sum_{k=0}^{n-1} \frac{(n-1)!}{k!((n-1)-k)!} p^k (1-p)^{(n-1)-k}\end{aligned}$$

Expectation of a Binomial R. V. - 2nd way

Using linearity of expectations

$$\mathbf{E}[X] = \mathbf{E} \left[\sum_{i=1}^n X_i \right] = \sum_{i=1}^n \mathbf{E}[X_i] = np.$$

Computing Expectation

Consider a discrete random variable X that takes values $1, 2, \dots, k$. Sometimes is it easier to use the following equation to compute the expectation.

$$\mathbf{E}[X] = \sum_{i=1}^k \mathbf{Pr}(X \geq i).$$

Proof?

Expectation is not everything....

Which Job Would You Prefer?

- A job that pays \$1000 a week.
- A job that pays \$1 a week plus a bonus of \$1,000,000 with probability $\frac{1}{1000}$.

Variance

Definition

The **variance** of a random variable X is

$$\text{Var}[X] = \mathbf{E}[(X - \mathbf{E}[X])^2] = \mathbf{E}[X^2] - (\mathbf{E}[X])^2.$$

Definition

The **standard deviation** of a random variable X is

$$\sigma(X) = \sqrt{\text{Var}[X]}.$$

Exercises

- **Question:** We have two events that are disjoint. Are they independent?

Exercises

- **Question:** We have two events that are disjoint. Are they independent?
- **Answer:** No. Clearly they are dependent. If one happens the probability of the other happening is zero.

Exercises

- **Question:** We have two events that are disjoint. Are they independent?
- **Answer:** No. Clearly they are dependent. If one happens the probability of the other happening is zero.
- **Question:** A coin has probability p of being head. What is the probability that I throw the coin 10 times and I get **all** heads?

Exercises

- **Question:** We have two events that are disjoint. Are they independent?
- **Answer:** No. Clearly they are dependent. If one happens the probability of the other happening is zero.
- **Question:** A coin has probability p of being head. What is the probability that I throw the coin 10 times and I get **all** heads?
- **Answer:** Each coin toss is independent. Therefore the probability is: p^{10} .

Exercises

- **Question:** We have two events that are disjoint. Are they independent?
- **Answer:** No. Clearly they are dependent. If one happens the probability of the other happening is zero.
- **Question:** A coin has probability p of being head. What is the probability that I throw the coin 10 times and I get **all** heads?
- **Answer:** Each coin toss is independent. Therefore the probability is: p^{10} .
- **Question:** A coin has probability p of being head. What is the probability that I throw the coin 10 times and I get **at least one** head?

Exercises

- **Question:** We have two events that are disjoint. Are they independent?
- **Answer:** No. Clearly they are dependent. If one happens the probability of the other happening is zero.
- **Question:** A coin has probability p of being head. What is the probability that I throw the coin 10 times and I get **all** heads?
- **Answer:** Each coin toss is independent. Therefore the probability is: p^{10} .
- **Question:** A coin has probability p of being head. What is the probability that I throw the coin 10 times and I get **at least one** head?
- **Answer:** Consider the case that I get no heads. Each coin toss is independent. Therefore the probability of getting no heads is $(1 - p)^{10}$.
- The probability of getting at least one head is $1 - (1 - p)^{10}$.

Exercise

- Assume that N people checked coats in a restaurants. The coats are mixed and each person gets a random coat.
- How many people we expect to have gotten their own coats?

Exercise

- Assume that N people checked coats in a restaurant. The coats are mixed and each person gets a random coat.
- How many people do we expect to have gotten their own coats?
- Let $X =$ “number of people that got their own coats”

Exercise

- Assume that N people checked coats in a restaurants. The coats are mixed and each person gets a random coat.
- How many people we expect to have gotten their own coats?
- Let $X =$ “number of people that got their own coats”
- It's hard to compute $E[X] = \sum_{k=0}^N k \Pr(X = k)$.

Exercise

- Assume that N people checked coats in a restaurants. The coats are mixed and each person gets a random coat.
- How many people we expect to have gotten their own coats?
- Let $X =$ “number of people that got their own coats”
- It's hard to compute $E[X] = \sum_{k=0}^N k \Pr(X = k)$.
- Instead we define N 0-1 random variables X_i :

$$X_i = \begin{cases} 1, & \text{if person } i \text{ got his coat,} \\ 0, & \text{otherwise} \end{cases}$$

Exercise

- Assume that N people checked coats in a restaurants. The coats are mixed and each person gets a random coat.
- How many people we expect to have gotten their own coats?
- Let $X =$ “number of people that got their own coats”
- It's hard to compute $E[X] = \sum_{k=0}^N k \Pr(X = k)$.
- Instead we define N 0-1 random variables X_i :

$$X_i = \begin{cases} 1, & \text{if person } i \text{ got his coat,} \\ 0, & \text{otherwise} \end{cases}$$

- $E[X_i] = 1 \cdot \Pr(X_i = 1) + 0 \cdot \Pr(X_i = 0) =$
- $\Pr(X_i = 1) = \frac{1}{N}$
- $E[X] = \sum_{i=1}^N E[X_i] = 1$

- What is the probability that everyone gets their own coat?

- What is the probability that everyone gets their own coat?
- **Incorrect argument:** The probability that one person gets their coat is $\Pr(X_i = 1) = 1/N$.
- The probability that everyone gets their coat is

$$\prod_{i=1}^N \Pr(X_i = 1) = \frac{1}{N^N}$$

Where is the error in this?

- What is the probability that everyone gets their own coat?
- **Incorrect argument:** The probability that one person gets their coat is $\Pr(X_i = 1) = 1/N$.
- The probability that everyone gets their coat is

$$\prod_{i=1}^N \Pr(X_i = 1) = \frac{1}{N^N}$$

Where is the error in this?

- The random variables are **not independent**. Once one person has found their coat the probability for the rest changes.
- What is the correct probability?

- What is the probability that everyone gets their own coat?
- **Incorrect argument:** The probability that one person gets their coat is $\Pr(X_i = 1) = 1/N$.
- The probability that everyone gets their coat is

$$\prod_{i=1}^N \Pr(X_i = 1) = \frac{1}{N^N}$$

Where is the error in this?

- The random variables are **not independent**. Once one person has found their coat the probability for the rest changes.
- What is the correct probability?
- One way to compute it:

$$\Pr(X_1)\Pr(X_2 | X_1) \cdots \Pr(X_N | X_{N-1}, \dots, X_1) = \frac{1}{N} \frac{1}{N-1} \cdots 1 = \frac{1}{N!}$$

- It also follows from the fact that of all possible permutations of coats there is only one that is the correct one.